

# Limits and Derivatives

---

## Limit of a Function Using Intuitive Approach

For a function  $f(x)$ , if for  $x$  closes to  $a$  implies that  $f(x)$  closes to  $l$ , then  $l$  is called the **limit** of function  $f(x)$  at  $a$ .

- $l$  is the limit of function  $f(x)$  is written as  $\lim_{x \rightarrow a} f(x) = l$  [read as “limit of  $f(x)$  is  $l$ , when  $x$  tends to  $a$ ” or “for  $x \rightarrow a$  ( $x$  tends to  $a$ ),  $f(x) \rightarrow l$  ( $f(x)$  tends to  $l$ )”]
- If  $f(x) = x^3 - 2$ , then for  $x$  very close to 3,  $f(x)$  will be very close to 25. This can be written as  $\lim_{x \rightarrow 3} (x^3 - 2) = 25$ . So, limiting value of  $x^3 - 2$  at  $x$  closes to 3 is 25.

**Example 1:** For  $f(x) = x(a - 3x)$ , find the value of  $a$  at which the limits of function  $f(x)$  when  $x$  tends to 4 and when it tends to 5 are the same?

**Solution:**

It is given that

$$f(x) = x(a - 3x)$$

$$\Rightarrow f(x) = ax - 3x^2$$

The limit of function  $f(x)$  when  $x$  tends to 4 is calculated as follows:

x	3.9	3.95	3.99	3.999	4.001	4.01	4.05	4.1
f(x)	3.9a - 45.63	3.95a - 46.8075	3.99a - 47.7603	3.999a - 47.976003	4.001a - 48.024003	4.01a - 48.2403	3.05a - 49.2075	4.1a - 50.43

$$\therefore \lim_{x \rightarrow 4} f(x) = \lim_{x \rightarrow 4} (ax - 3x^2) = 4a - 48$$

The limit of function  $f(x)$  when  $x$  tends to 5 is calculated as follows:

x	4.9	4.95	4.99	4.999	5.001	5.01	5.05	5.1
---	-----	------	------	-------	-------	------	------	-----

$f(x)$	$4.9a - 7$	$4.95a - 7$	$4.99a - 7$	$4.999a - 7$	$5.001a - 7$	$5.01a - 7$	$5.05a - 7$	$5.1a - 7$
	2.03	3.5075	4.7003	970003	3	3003	5075	8.03

$$\therefore \lim_{x \rightarrow 5} f(x) = \lim_{x \rightarrow 5} (ax - 3x^2) = 5a - 75$$

We have to find the particular value of  $a$  at which the limits of function  $f(x)$  when  $x$  tends to 4 and when it tends to 5 are equal.

$$\begin{aligned} \therefore \lim_{x \rightarrow 4} f(x) &= \lim_{x \rightarrow 5} f(x) \\ \Rightarrow 4a - 48 &= 5a - 75 \\ \Rightarrow a &= 27 \end{aligned}$$

Thus, the limiting values of  $f(x) = x(a - 3x)$  when  $x$  tends to 4 and 5 are equal for  $a = 27$ .

**Example 2:** Show that the limit value of  $g(y) = [2y - 5]$  does not exist when  $y$  tends to 2.

**Solution:** The given function is

$$g(y) = [2y - 5].$$

Clearly,  $g(y)$  is a greatest integer function

$$\text{Hence, } g(y) = \begin{cases} a-1, & \text{for } a-1 < g(y) < a \\ a, & \text{for } a \leq g(y) < a+1 \end{cases}$$

Where,  $a$  is an integer

The limit of  $g(y)$  when  $y$  tends to 2 is calculated as follows:

$y$	1.9	1.95	1.99	1.999	2.001	2.01	2.05	2.1
$g(y)$	-2	-2	-2	-2	-1	-1	-1	-1

We may observe that

Left hand limit of the function =  $\lim_{y \rightarrow 2^-} g(y) = -2$  whereas the right hand limit =  $\lim_{y \rightarrow 2^+} g(y) = -1$ .

Since the left hand and the right hand limits of the function are not equal, the given function does not have a limiting value.

**Example 3:** For what real and complex values of  $b$ ,  $\lim_{t \rightarrow b} v(t) \neq v(b)$ ,

where 
$$v(t) = \frac{(t^4 - 16)(t^2 - 16)}{(t^3 - 1)(2t^2 - t - 28)}$$
 ?

**Solution:**

We know that if a function  $v(t)$  is defined at  $t = b$ , then  $\lim_{t \rightarrow b} v(t) = v(b)$ , else not.

Since  $\lim_{t \rightarrow b} v(t) \neq v(b)$ , we need to find the value of  $b$ , i.e.,  $t$ , where  $v(t)$  does not exist.

This is only possible, if

$$\begin{aligned} (t^3 - 1)(2t^2 - t - 28) &= 0 \\ \Rightarrow (t - 1)(t^2 + t + 1)(2t^2 - 8t + 7t - 28) &= 0 \\ \Rightarrow (t - 1)(t^2 + t + 1)[2t(t - 4) + 7(t - 4)] &= 0 \\ \Rightarrow (t - 1)(t^2 + t + 1)(t - 4)(2t + 7) &= 0 \\ \Rightarrow t = 1 \text{ or } 4 \text{ or } \frac{-7}{2} \text{ or } \frac{-1 \pm \sqrt{1^2 - 4(1)(1)}}{2(1)} \\ \Rightarrow t = 1 \text{ or } 4 \text{ or } \frac{-7}{2} \text{ or } \frac{-1 \pm i\sqrt{3}}{2} \end{aligned}$$

So, for  $b = 1, 4, \frac{-7}{2}$  as real values and  $b = \frac{-1 \pm i\sqrt{3}}{2}$  as the complex values,  $\lim_{t \rightarrow b} v(t) \neq v(b)$

, where 
$$v(t) = \frac{(t^4 - 16)(t^2 - 16)}{(t^3 - 1)(2t^2 - t - 28)}$$
.

### Limit of a Polynomial and a Rational Function

## Algebra of Limits

- If  $f$  and  $g$  are two functions such that both  $\lim_{x \rightarrow a} f(x)$  and  $\lim_{x \rightarrow a} g(x)$  exist, then

$$\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

The limit of the sum of two functions is the sum of the limits of the functions.

$$\lim_{x \rightarrow 4} \left( x^{\frac{5}{2}} + x^{\frac{3}{2}} \right) = \lim_{x \rightarrow 4} x^{\frac{5}{2}} + \lim_{x \rightarrow 4} x^{\frac{3}{2}} = 4^{\frac{5}{2}} + 4^{\frac{3}{2}} = 32 + 8 = 40$$

For example,

$$\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

The limit of the difference between two functions is the difference between the limits of the functions.

$$\lim_{x \rightarrow 4} \left( x^{\frac{5}{2}} - x^{\frac{3}{2}} \right) = \lim_{x \rightarrow 4} x^{\frac{5}{2}} - \lim_{x \rightarrow 4} x^{\frac{3}{2}} = 4^{\frac{5}{2}} - 4^{\frac{3}{2}} = 32 - 8 = 24$$

For example,

$$\lim_{x \rightarrow a} [f(x).g(x)] = \lim_{x \rightarrow a} f(x). \lim_{x \rightarrow a} g(x)$$

The limit of the product of two functions is the product of the limits of the functions.

$$\lim_{x \rightarrow 4} \left( x^{\frac{5}{2}}.x^{\frac{3}{2}} \right) = \lim_{x \rightarrow 4} x^{\frac{5}{2}}. \lim_{x \rightarrow 4} x^{\frac{3}{2}} = 4^{\frac{5}{2}} \times 4^{\frac{3}{2}} = 32 \times 8 = 256$$

For example,

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \text{ where } \lim_{x \rightarrow a} g(x) \neq 0$$

The limit of the quotient of the two functions is the quotient of the limits of the functions, where the denominator is not zero.

$$\lim_{x \rightarrow 4} \frac{x^{\frac{5}{2}}}{x^{\frac{3}{2}}} = \frac{\lim_{x \rightarrow 4} x^{\frac{5}{2}}}{\lim_{x \rightarrow 4} x^{\frac{3}{2}}} = \frac{4^{\frac{5}{2}}}{4^{\frac{3}{2}}} = \frac{32}{8} = 4$$

For example,

$$\lim_{x \rightarrow a} [k.f(x)] = k \lim_{x \rightarrow a} f(x), \text{ where } k \text{ is a constant}$$

The limit of the product of a constant and a function is the product of the constant and the limit of that function.

$$\lim_{x \rightarrow 4} \left( \frac{9}{2} x^{\frac{5}{2}} \right) = \frac{9}{2} \lim_{x \rightarrow 4} x^{\frac{5}{2}} = \frac{9}{2} \times 4^{\frac{5}{2}} = \frac{9}{2} \times 32 = 144$$

For example,

## Limit of a Polynomial Function

- A function  $p(x)$  is said to be a polynomial function if  $p(x) = 0$  or  $p(x) = \sum_{r=0}^n a_r x^r$ , where  $a_r \in \mathbb{R}$  and  $a_r \neq 0$  for some whole number  $r$ .
- The limit of a polynomial function  $p(x)$  at  $x = a$  is given by  $\lim_{x \rightarrow a} p(x) = p(a)$

For example, the value of  $\lim_{m \rightarrow n+3} (3m^3 - 9m^2n + 9mn^2 - 3n^3 - m + n - 80)$  can be calculated as follows:

$$\begin{aligned} & \lim_{m \rightarrow n+3} (3m^3 - 9m^2n + 9mn^2 - 3n^3 - m + n - 80) \\ &= \lim_{m \rightarrow n+3} [3(m^3 - 3m^2n + 3mn^2 - n^3) - (m - n) - 80] \\ &= \lim_{m \rightarrow n+3} [3(m - n)^3 - (m - n) - 80] \\ &= [3(3)^3 - (3) - 80] \\ &= 81 - 3 - 80 \\ &= -2 \end{aligned}$$

### Limit of a Rational Function

- A function  $p(x)$  is said to be a rational function if  $p(x) = \frac{q(x)}{r(x)}$ , where  $q(x)$  and  $r(x)$  are polynomials such that  $r(x) \neq 0$ .
- The limit of a rational function  $p(x)$  of the form  $p(x) = \frac{q(x)}{r(x)}$  at  $x = a$  is given by

$$\lim_{x \rightarrow a} p(x) = \frac{q(a)}{r(a)}$$

- For example, to find the value of  $\lim_{x \rightarrow 64} \frac{\sqrt{x} + 7}{\sqrt[3]{x} + 2}$ , we may proceed as follows.

$$\lim_{x \rightarrow 64} \frac{\sqrt{x} + 7}{\sqrt[3]{x} + 2} = \frac{\sqrt{64} + 7}{\sqrt[3]{64} + 2} = \frac{8 + 7}{4 + 2} = \frac{15}{6} = \frac{5}{2}$$

- For any positive integer  $n$ ,  $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$

- For example,  $\lim_{y \rightarrow 0} \frac{(y+5)^4 - 625}{y}$  can be calculated as follows.

$$\begin{aligned} \lim_{y \rightarrow 0} \frac{(y+5)^4 - 625}{y} &= \lim_{y+5 \rightarrow 5} \frac{(y+5)^4 - 5^4}{(y+5) - 5} && (y \rightarrow 0 \text{ shows that } y+5 \rightarrow 5) \\ &= 4 \times 5^{4-1} \\ &= 500 \end{aligned}$$

**Example 1:** Find the values of  $a$  and  $b$  if

$$\lim_{n \rightarrow \infty} \frac{3a.(n+5)! - 2b.(n+4)!}{b.(n+5)! + a.(n+4)!} = -2 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{(a+2b).(n+1)! - b.(n-1)!}{(2a-b+1).(n+1)! - a.(n-1)!} = \frac{-1}{2}$$

Also, show that 
$$\lim_{x \rightarrow 1} \frac{a+2b}{x} = \lim_{x \rightarrow \frac{-3}{2}} \frac{b-a}{x^2-1}$$
.

**Solution:**

We have 
$$\lim_{n \rightarrow \infty} \frac{3a.(n+5)! - 2b.(n+4)!}{b.(n+5)! + a.(n+4)!} = -2$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{[3a.(n+5) - 2b](n+4)!}{[b.(n+5) + a](n+4)!} = -2$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{3an + 15a - 2b}{bn + 5b + a} = -2$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n \left( 3a + \frac{15a - 2b}{n} \right)}{n \left( b + \frac{5b + a}{n} \right)} = -2$$

$$\Rightarrow \frac{\lim_{n \rightarrow \infty} \left( 3a + \frac{15a - 2b}{n} \right)}{\lim_{n \rightarrow \infty} \left( b + \frac{5b + a}{n} \right)} = -2$$

$$\Rightarrow \frac{3a + 0}{b + 0} = -2$$

$$\Rightarrow 3a = -2b$$

$$\Rightarrow a = \frac{-2b}{3}$$

... (1)

We also have 
$$\lim_{n \rightarrow \infty} \frac{(a+2b).(n+1)! - b.(n-1)!}{(2a-b+1).(n+1)! - a.(n-1)!} = \frac{-1}{2}$$

$$\begin{aligned}
&\Rightarrow \lim_{n \rightarrow \infty} \frac{[(a+2b)n(n+1)-b](n-1)!}{[(2a-b+1)n(n+1)-a](n-1)!} = \frac{-1}{2} \\
&\Rightarrow \lim_{n \rightarrow \infty} \frac{(a+2b)n^2 + (a+2b)n - b}{(2a-b+1)n^2 + (2a-b+1)n - a} = \frac{-1}{2} \\
&\Rightarrow \lim_{n \rightarrow \infty} \frac{n^2 \left[ (a+2b) + \frac{(a+2b)}{n} - \frac{b}{n^2} \right]}{n^2 \left[ (2a-b+1) + \frac{(2a-b+1)}{n} - \frac{a}{n^2} \right]} = \frac{-1}{2} \\
&\Rightarrow \frac{\lim_{n \rightarrow \infty} \left[ (a+2b) + \frac{(a+2b)}{n} - \frac{b}{n^2} \right]}{\lim_{n \rightarrow \infty} \left[ (2a-b+1) + \frac{(2a-b+1)}{n} - \frac{a}{n^2} \right]} = \frac{-1}{2} \\
&\Rightarrow \frac{a+2b}{2a-b+1} = \frac{-1}{2} \\
&\Rightarrow \frac{\frac{-2b}{3} + 2b}{2 \times \frac{-2b}{3} - b + 1} = \frac{-1}{2} \quad \text{[Using equation (1)]} \\
&\Rightarrow \frac{4b}{-7b+3} = \frac{-1}{2} \\
&\Rightarrow 8b = 7b - 3 \\
&\Rightarrow b = -3
\end{aligned}$$

Substituting the value of  $b$  in equation (1), we obtain

$$a = 2$$

Hence,  $a = 2$  and  $b = -3$

Now,

$$\lim_{x \rightarrow 1} \frac{a+2b}{x} = \frac{2+2(-3)}{1} = -4$$

$$\lim_{x \rightarrow \frac{-3}{2}} \frac{b-a}{x^2-1} = \frac{(-3)-2}{\left(\frac{-3}{2}\right)^2-1} = \frac{-5}{\frac{5}{4}} = -4$$

and

$$\Rightarrow \Rightarrow \lim_{x \rightarrow 1} \frac{a+2b}{x} = \lim_{x \rightarrow \frac{-3}{2}} \frac{b-a}{x^2-1}$$

**Example 2:** Find the value of  $n$ , such that  $\lim_{a \rightarrow b-3} \frac{(a-b)^{2n}-9^n}{(a-b)^{3n}+27^n} = -\frac{2}{729}$ , where  $n$  is an odd number.

**Solution:**

$$\lim_{a \rightarrow b-3} \frac{(a-b)^{2n}-9^n}{(a-b)^{3n}+27^n} = -\frac{2}{729}$$

$$\Rightarrow \lim_{a \rightarrow b-3} \frac{(a-b)^{2n}-3^{2n}}{(a-b)^{3n}+3^{3n}} = -\frac{2}{729} \quad (a \rightarrow b-3 \Rightarrow a-b \rightarrow -3)$$

$$\Rightarrow \lim_{a \rightarrow b-3} \frac{(a-b)^{2n}-(-3)^{2n}}{(a-b)^{3n}-(-3)^{3n}} = -\frac{2}{729} \quad (\text{Since } n \text{ is an odd number, } (-3)^{2n} = 3^{2n} \text{ and } (-3)^{3n} = -3^{3n})$$

$$\Rightarrow \frac{\lim_{a \rightarrow b-3} \frac{(a-b)^{2n}-(-3)^{2n}}{(a-b)-(-3)}}{\lim_{a \rightarrow b-3} \frac{(a-b)^{3n}-(-3)^{3n}}{(a-b)-(-3)}} = -\frac{2}{729}$$

$$\Rightarrow \frac{2n(-3)^{2n-1}}{3n(-3)^{3n-1}} = -\frac{2}{729}$$

$$\Rightarrow \frac{2}{3(-3)^n} = -\frac{2}{729}$$

$$\Rightarrow (-3)^n = \frac{-729}{3}$$

$$\Rightarrow (-3)^n = -243 = (-3)^5$$

$$\Rightarrow n = 5$$

**Example 3:** Evaluate  $\lim_{x \rightarrow 0} \frac{\sqrt{4+x^3} - \sqrt{4+x}}{\sqrt{9+x^7} - \sqrt{9+x}}$

**Solution:**

$$\lim_{x \rightarrow 0} \frac{\sqrt{4+x^3} - \sqrt{4+x}}{\sqrt{9+x^7} - \sqrt{9+x}} = \frac{0}{0} \text{ form}$$

Hence,

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{\sqrt{4+x^3} - \sqrt{4+x}}{\sqrt{9+x^7} - \sqrt{9+x}} \\ &= \lim_{x \rightarrow 0} \left( \left( \sqrt{4+x^3} - \sqrt{4+x} \right) \times \frac{1}{\sqrt{9+x^7} - \sqrt{9+x}} \right) \\ &= \lim_{x \rightarrow 0} \left( \frac{\left( \sqrt{4+x^3} - \sqrt{4+x} \right) \left( \sqrt{4+x^3} + \sqrt{4+x} \right)}{\sqrt{4+x^3} + \sqrt{4+x}} \times \frac{\sqrt{9+x^7} + \sqrt{9+x}}{\left( \sqrt{9+x^7} - \sqrt{9+x} \right) \left( \sqrt{9+x^7} + \sqrt{9+x} \right)} \right) \\ &= \lim_{x \rightarrow 0} \left( \frac{(4+x^3) - (4+x)}{\sqrt{4+x^3} + \sqrt{4+x}} \times \frac{\sqrt{9+x^7} + \sqrt{9+x}}{(9+x^7) - (9+x)} \right) \\ &= \lim_{x \rightarrow 0} \left( \frac{x(x^2-1)}{\sqrt{4+x^3} + \sqrt{4+x}} \times \frac{\sqrt{9+x^7} + \sqrt{9+x}}{x(x^6-1)} \right) \\ &= \lim_{x \rightarrow 0} \left( \frac{x^2-1}{x^6-1} \times \frac{\sqrt{9+x^7} + \sqrt{9+x}}{\sqrt{4+x^3} + \sqrt{4+x}} \right) \\ &= \lim_{x \rightarrow 0} \frac{x^2-1}{x^6-1} \times \lim_{x \rightarrow 0} \frac{\sqrt{9+x^7} + \sqrt{9+x}}{\sqrt{4+x^3} + \sqrt{4+x}} \\ &= \frac{-1}{-1} \times \frac{3+3}{2+2} \\ &= \frac{3}{2} \end{aligned}$$

### Limits of Trigonometric Functions

- Let  $f$  and  $g$  be two real-valued functions with the same domain, such that  $f(x) \leq g(x)$  for all  $x$  in the domain of definition. For some  $a$ , if both  $\lim_{x \rightarrow a} f(x)$  and  $\lim_{x \rightarrow a} g(x)$  exist, then  $\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x)$ .
- For example, we know that  $x^2 \leq x^3$ , for  $x \in \mathbf{R}$  and  $x \geq 1$ . So, for any  $a \in \mathbf{R}$  and  $a \geq 1$ ,  $\lim_{x \rightarrow a} x^2 \leq \lim_{x \rightarrow a} x^3$ .
- Two important limits are

- $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

- $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$

$$\lim_{x \rightarrow \frac{\pi}{3}} \frac{\sqrt{3} \sin\left(\frac{\pi}{2} - x\right) + \sin(\pi + x)}{3\pi\left(\frac{\pi}{3} - x\right)}$$

**Example 1:** Evaluate

**Solution**

$$\begin{aligned} \lim_{x \rightarrow \frac{\pi}{3}} \frac{\sqrt{3} \sin\left(\frac{\pi}{2} - x\right) + \sin(\pi + x)}{3\pi\left(\frac{\pi}{3} - x\right)} &= \lim_{\frac{\pi}{3} - x \rightarrow 0} \frac{\sqrt{3} \cos x - \sin x}{3\pi\left(\frac{\pi}{3} - x\right)} \\ &= \frac{1}{3\pi} \cdot \lim_{\frac{\pi}{3} - x \rightarrow 0} \frac{2 \cdot \left[ \frac{\sqrt{3}}{2} \cos x - \frac{1}{2} \sin x \right]}{\frac{\pi}{3} - x} \\ &= \frac{2}{3\pi} \cdot \lim_{\frac{\pi}{3} - x \rightarrow 0} \frac{\left[ \sin \frac{\pi}{3} \cos x - \cos \frac{\pi}{3} \sin x \right]}{\frac{\pi}{3} - x} \\ &= \frac{2}{3\pi} \cdot \lim_{\frac{\pi}{3} - x \rightarrow 0} \frac{\sin\left(\frac{\pi}{3} - x\right)}{\frac{\pi}{3} - x} \\ &= \frac{2}{3\pi} \times 1 \\ &= \frac{2}{3\pi} \end{aligned}$$

**Example 2:** If  $\lim_{x \rightarrow 0} \frac{\cos 4x - \sin\left(\frac{\pi}{2} + 5x\right)}{x^2} = \frac{3a+b}{2}$  and

$\lim_{x \rightarrow 0} \frac{\sin\left(\frac{\pi}{4} + 5x\right) - \sin\left(\frac{\pi}{4} + 3x\right)}{x} = \sqrt{4b-5a}$ , then find the value of  $\sqrt{5a+2b}$ .

**Solution:**

$$\lim_{x \rightarrow 0} \frac{\cos 4x - \sin\left(\frac{\pi}{2} + 5x\right)}{x^2} = \frac{3a+b}{2}$$

$$\begin{aligned} \Rightarrow \frac{3a+b}{2} &= \lim_{x \rightarrow 0} \frac{\cos 4x - \cos 5x}{x^2} \\ &= \lim_{x \rightarrow 0} \frac{2 \sin\left(\frac{5x+4x}{2}\right) \sin\left(\frac{5x-4x}{2}\right)}{x^2} \\ &= 2 \lim_{x \rightarrow 0} \frac{\sin \frac{9x}{2} \cdot \sin \frac{x}{2}}{x^2} \\ &= 2 \lim_{x \rightarrow 0} \frac{\sin \frac{9x}{2}}{x} \times \lim_{x \rightarrow 0} \frac{\sin \frac{x}{2}}{x} \\ &= 2 \times \frac{9}{2} \lim_{\frac{9x}{2} \rightarrow 0} \frac{\sin \frac{9x}{2}}{\frac{9x}{2}} \times \frac{1}{2} \cdot \lim_{\frac{x}{2} \rightarrow 0} \frac{\sin \frac{x}{2}}{\frac{x}{2}} \\ &= 9 \times 1 \times \frac{1}{2} \times 1 \\ &= \frac{9}{2} \\ \Rightarrow 3a + b &= 9 \\ \Rightarrow b &= 9 - 3a \dots(1) \end{aligned}$$

It is also given that

$$\lim_{x \rightarrow 0} \frac{\sin\left(\frac{\pi}{4} + 5x\right) - \sin\left(\frac{\pi}{4} + 3x\right)}{x} = \sqrt{4b - 5a}$$

$$\Rightarrow \sqrt{4b - 5a} = \lim_{x \rightarrow 0} \frac{2 \sin\left[\frac{\left(\frac{\pi}{4} + 5x\right) - \left(\frac{\pi}{4} + 3x\right)}{2}\right] \cdot \cos\left[\frac{\left(\frac{\pi}{4} + 5x\right) + \left(\frac{\pi}{4} + 3x\right)}{2}\right]}{x}$$

$$= 2 \lim_{x \rightarrow 0} \frac{\sin x \cdot \cos\left(\frac{\pi}{4} + 4x\right)}{x}$$

$$= 2 \lim_{x \rightarrow 0} \frac{\sin x}{x} \cdot \lim_{x \rightarrow 0} \cos\left(\frac{\pi}{4} + 4x\right)$$

$$= 2 \times 1 \times \frac{1}{\sqrt{2}}$$

$$= \sqrt{2}$$

$$\Rightarrow 4b - 5a = 2$$

From (1), we have

$$4(9 - 3a) - 5a = 2$$

$$36 - 17a = 2$$

$$17a = 34$$

$$a = 2$$

Substituting  $a = 2$  in equation (1), we obtain  $b = 3$

$$\text{Now, } \sqrt{5a + 2b} = \sqrt{5 \times 2 + 2 \times 3} = \sqrt{16} = 4$$

### Derivative of a Function

- Suppose  $f$  is a real-valued function and  $a$  is a point in the domain of definition. If the

limit  $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$  exists, then it is called the derivative of  $f$  at  $a$ . The derivative

of  $f$  at  $a$  is denoted by  $f'(a)$ .

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

- Suppose  $f$  is a real-valued function. The derivative of  $f$  {denoted by  $f'(x)$  or  $\frac{d}{dx}[f(x)]$ } is defined by

$$\frac{d}{dx}[f(x)] = f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

This definition of derivative is called the **first principle** of derivative.

- For example, the derivative of  $y = (ax - b)^{10}$  is calculated as follows.

We have  $y = f(x) = (ax - b)^{10}$ ; using the first principle of derivative, we obtain

$$\begin{aligned} \frac{dy}{dx} = f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[a(x+h) - b]^{10} - (ax - b)^{10}}{h} \\ &= \lim_{h \rightarrow 0} \frac{[a(x+h) - b - (ax - b)] \cdot \sum_{r=0}^9 [a(x+h) - b]^{9-r} (ax - b)^r}{h} \\ &= \lim_{h \rightarrow 0} \frac{ah}{h} \cdot \lim_{h \rightarrow 0} \sum_{r=0}^9 [a(x+h) - b]^{9-r} (ax - b)^r \\ &= a \sum_{r=0}^9 (ax - b)^{9-r} \cdot (ax - b)^r \\ &= a[(ax - b)^{9-0} \cdot (ax - b)^0 + (ax - b)^{9-1} \cdot (ax - b)^1 + \dots + (ax - b)^{9-9} \cdot (ax - b)^9] \\ &= 10a(ax - b)^9 \end{aligned}$$

### Solved Examples

**Example 1:** Find the derivative of  $f(x) = \operatorname{cosec}^2 2x + \tan^2 4x$ . Also, find  $f'(x)$  at  $x = \frac{\pi}{6}$ .

**Solution:**

The derivative of  $f(x) = \operatorname{cosec}^2 2x + \tan^2 4x$  is calculated as follows.

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{\operatorname{cosec}^2 2(x+h) + \tan^2 4(x+h) - [\operatorname{cosec}^2 2(x) + \tan^2 4(x)]}{h} \\
&= \lim_{h \rightarrow 0} \frac{[\operatorname{cosec}^2(2x+2h) - \operatorname{cosec}^2 2x] + [\tan^2(4x+4h) - \tan^2(4x)]}{h} \\
&= \lim_{h \rightarrow 0} \frac{\left( \frac{1}{\sin^2(2x+2h)} - \frac{1}{\sin^2 2x} \right) + \left( \frac{\sin^2(4x+4h)}{\cos^2(4x+4h)} - \frac{\sin^2 4x}{\cos^2 4x} \right)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\left( \frac{\sin^2 2x - \sin^2(2x+2h)}{\sin^2 2x \sin^2(2x+2h)} \right) + \left( \frac{\sin^2(4x+4h)\cos^2 4x - \cos^2(4x+4h)\sin^2 4x}{\cos^2 4x \cos^2(4x+4h)} \right)}{h} \\
&= \lim_{h \rightarrow 0} \frac{[\sin 2x - \sin(2x+2h)][\sin 2x + \sin(2x+2h)]}{h \sin^2 2x \sin^2(2x+2h)} \\
&\quad + \lim_{h \rightarrow 0} \frac{[\sin(4x+4h)\cos 4x - \cos(4x+4h)\sin 4x][\sin(4x+4h)\cos 4x + \cos(4x+4h)\sin 4x]}{h \cos^2 4x \cos^2(4x+4h)} \\
&= \lim_{h \rightarrow 0} \frac{2 \cos(2x+h) \sin(-h) \times 2 \sin(2x+h) \cos(-h)}{h \sin^2 2x \sin^2(2x+2h)} + \lim_{h \rightarrow 0} \frac{\sin(4x+4h-4x)\sin(4x+4h+4x)}{h \cos^2 4x \cos^2(4x+4h)} \\
&= -4 \lim_{h \rightarrow 0} \frac{\sin h}{h} \times \lim_{h \rightarrow 0} \frac{\cos(2x+h) \times \sin(2x+h) \cos(h)}{\sin^2 2x \sin^2(2x+2h)} + 4 \lim_{h \rightarrow 0} \frac{\sin(4h)}{4h} \times \lim_{h \rightarrow 0} \frac{\sin(4x+4h+4x)}{\cos^2 4x \cos^2(4x+4h)} \\
&= -4 \times 1 \times \frac{\cos 2x}{\sin^3 2x} + 4 \times 1 \times \frac{\sin 8x}{\cos^4 4x} \\
&= -4 \cot 2x \operatorname{cosec}^2 2x + \frac{8 \sin 4x \cos 4x}{\cos^4 4x} \\
&= -4 \cot 2x \operatorname{cosec}^2 2x + 8 \tan 4x \sec^2 4x
\end{aligned}$$

At  $x = \frac{\pi}{6}$ ,  $f'\left(\frac{\pi}{6}\right)$  is given by

$$\begin{aligned}
f'\left(\frac{\pi}{6}\right) &= -4 \cot\left(\frac{\pi}{3}\right) \operatorname{cosec}^2\left(\frac{\pi}{3}\right) + 8 \tan\left(\frac{2\pi}{3}\right) \sec^2\left(\frac{2\pi}{3}\right) \\
&= -4 \times \frac{1}{\sqrt{3}} \times \left(\frac{2}{\sqrt{3}}\right)^2 + 8(-\sqrt{3}) \times (-2)^2 \\
&= \frac{-16}{3\sqrt{3}} - 32\sqrt{3} \\
&= \frac{-304}{3\sqrt{3}}
\end{aligned}$$

**Example 2:** If  $y = (ax^2 + x + b)^2$ , then find the values of  $a$  and  $b$ , such

that  $\frac{dy}{dx} = 4x^2(4x+3) + 2(13x+3)$ .

**Solution:**

It is given that  $y = (ax^2 + x + b)^2$

$$\begin{aligned} \Rightarrow \frac{dy}{dx} &= \lim_{h \rightarrow 0} \frac{[a(x+h)^2 + (x+h) + b]^2 - [ax^2 + x + b]^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{[a(x+h)^2 + (x+h) + b - (ax^2 + x + b)][a(x+h)^2 + (x+h) + b + (ax^2 + x + b)]}{h} \\ &= \lim_{h \rightarrow 0} \frac{[a(2xh + h^2) + h][a(x+h)^2 + (x+h) + b + (ax^2 + x + b)]}{h} \\ &= \lim_{h \rightarrow 0} \frac{h[a(2x+h) + 1]}{h} \times \lim_{h \rightarrow 0} [a(x+h)^2 + (x+h) + b + (ax^2 + x + b)] \\ &= (2ax + 1) \times 2(ax^2 + x + b) \\ &= 4a^2x^3 + 6ax^2 + (4ab + 2)x + 2b \\ \Rightarrow 4x^2(4x+3) + 2(13x+3) &= 4a^2x^3 + 6ax^2 + (4ab + 2)x + 2b \\ \Rightarrow 4a^2x^3 + 6ax^2 + (4ab + 2)x + 2b &= 16x^3 + 12x^2 + 26x + 6 \end{aligned}$$

Comparing the coefficients of  $x^3$ ,  $x^2$ ,  $x$ , and the constant terms of the above expression, we obtain

$$\begin{aligned} 4a^2 &= 16, \quad 6a = 12, \quad 4ab + 2 = 26 \text{ and } 2b = 6 \\ \Rightarrow a &= \pm 2, \quad a = 2, \quad b = 3 \text{ and } b = 3 \\ \Rightarrow a &= 2 \text{ and } b = 3 \end{aligned}$$

**Example 3:** What is the derivative of  $y$  with respect to  $x$ , if  $y = \sqrt{\frac{ax+b}{cx-d}}$  ?

**Solution:**

It is given that  $y = \sqrt{\frac{ax+b}{cx-d}}$

$$\begin{aligned}
\Rightarrow \frac{dy}{dx} &= \frac{d}{dx} \left( \sqrt{\frac{ax+b}{cx-d}} \right) \\
&= \lim_{h \rightarrow 0} \frac{\sqrt{\frac{a(x+h)+b}{c(x+h)-d}} - \sqrt{\frac{ax+b}{cx-d}}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sqrt{[a(x+h)+b](cx-d)} - \sqrt{[c(x+h)-d][ax+b]}}{h\sqrt{[c(x+h)-d][cx-d]}} \\
&\quad \left( \sqrt{[a(x+h)+b](cx-d)} - \sqrt{[c(x+h)-d][ax+b]} \right) \times \\
&= \lim_{h \rightarrow 0} \frac{\left( \sqrt{[a(x+h)+b](cx-d)} + \sqrt{[c(x+h)-d][ax+b]} \right)}{h\left( \sqrt{[c(x+h)-d][cx-d]} \right) \left( \sqrt{[a(x+h)+b](cx-d)} + \sqrt{[c(x+h)-d][ax+b]} \right)} \\
&= \lim_{h \rightarrow 0} \frac{[a(x+h)+b](cx-d) - [c(x+h)-d][ax+b]}{h\left( \sqrt{[c(x+h)-d][cx-d]} \right) \left( \sqrt{[a(x+h)+b](cx-d)} + \sqrt{[c(x+h)-d][ax+b]} \right)} \\
&= \lim_{h \rightarrow 0} \frac{h[a(cx-d) - c(ax+b)]}{h\left( \sqrt{[c(x+h)-d][cx-d]} \right) \left( \sqrt{[a(x+h)+b](cx-d)} + \sqrt{[c(x+h)-d][ax+b]} \right)} \\
&= \frac{a(cx-d) - c(ax+b)}{\left( \sqrt{[cx-d][cx-d]} \right) \left( \sqrt{(ax+b)(cx-d)} + \sqrt{(cx-d)(ax+b)} \right)} \\
&= \frac{-(ad+bc)}{2(cx-d)\sqrt{(ax+b)(cx-d)}}
\end{aligned}$$

## Derivatives of Trigonometric and Polynomial Functions

### Derivatives of Trigonometric Functions and Standard Formulas

- $\frac{d}{dx}(\sin x) = \cos x$

- $\frac{d}{dx}(\cos x) = -\sin x$

- $\frac{d}{dx}(x^n) = nx^{n-1}$

For example,  $\frac{d}{dx}(x^7) = 7x^{7-1} = 7x^6$

- $\frac{d}{dx}(C) = 0$ , where C is a constant

### Algebra of Derivatives

- If  $f$  and  $g$  are two functions such that their derivatives are defined in a common domain, then

- $$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

This means that the derivative of the sum of two functions is the sum of the derivatives of the functions.

For example, 
$$\frac{d}{dx}\left(x^{\frac{5}{2}} + x^{\frac{3}{2}}\right) = \frac{d}{dx}\left(x^{\frac{5}{2}}\right) + \frac{d}{dx}\left(x^{\frac{3}{2}}\right) = \frac{5}{2}x^{\frac{5}{2}-1} + \frac{3}{2}x^{\frac{3}{2}-1} = \frac{5}{2}x^{\frac{3}{2}} + \frac{3}{2}x^{\frac{1}{2}}$$

- $$\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

This means that the derivative of the difference between two functions is the difference between the derivatives of the function.

For example, 
$$\frac{d}{dx}\left(\sin x - x^{\frac{1}{3}}\right) = \frac{d}{dx}(\sin x) - \frac{d}{dx}\left(x^{\frac{1}{3}}\right) = \cos x - \frac{1}{3}x^{\frac{1}{3}-1} = \cos x - \frac{1}{3}x^{-\frac{2}{3}}$$

- $$\frac{d}{dx}[f(x).g(x)] = \frac{d}{dx}f(x).g(x) + f(x).\frac{d}{dx}g(x)$$

This is known as the **product** rule of derivative.

For example,

$$\frac{d}{dx}(x^3 \cos x) = \frac{d}{dx}(x^3).\cos x + (x^3).\frac{d}{dx}(\cos x) = 3x^2 \cos x + x^3(-\sin x) = 3x^2 \cos x - x^3 \sin x$$

- $$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{\frac{d}{dx}f(x).g(x) - f(x).\frac{d}{dx}g(x)}{[g(x)]^2}, \text{ where } \frac{d}{dx}g(x) \neq 0$$

This is known as the **quotient rule** of derivative.

For example,

$$\begin{aligned}
 \frac{d}{dx}(\tan x) &= \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) \\
 &= \frac{\frac{d}{dx}(\sin x) \cdot \cos x - \sin x \cdot \frac{d}{dx}(\cos x)}{(\cos x)^2} \\
 &= \frac{\cos x \cdot \cos x - \sin x(-\sin x)}{\cos^2 x} \\
 &= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\
 &= \frac{1}{\cos^2 x} \\
 &= \sec^2 x
 \end{aligned}$$

- $\frac{d}{dx}[k \cdot f(x)] = k \frac{d}{dx} f(x)$ , where  $k$  is a constant

This means that the derivative of the product of a constant and a function is the product of that constant and the derivative of that function.

For example,

$$\begin{aligned}
 \frac{d}{dx}(\sin 2x) &= \frac{d}{dx}(2 \sin x \cdot \cos x) \\
 &= 2 \frac{d}{dx}(\sin x \cdot \cos x) \\
 &= 2 \left( \frac{d}{dx}(\sin x) \cdot \cos x + \sin x \cdot \frac{d}{dx}(\cos x) \right) \\
 &= 2[\cos x \cdot \cos x + \sin x \cdot (-\sin x)] \\
 &= 2(\cos^2 x - \sin^2 x) \\
 &= 2 \cos 2x
 \end{aligned}$$

### Derivative of a Polynomial Function

- A function  $p(x)$  is said to be a polynomial function if  $p(x) = 0$  or  $p(x) = \sum_{r=0}^n a_r x^r$ , where  $a_r \in \mathbb{R}$  and  $a_r \neq 0$  for some whole number  $r$ .

- The derivative of a polynomial function  $p(x) = \sum_{r=0}^n a_r x^r$  is given by

$$\frac{d}{dx}[p(x)] = \sum_{r=1}^n r a_r x^{r-1}$$

## Solved Examples

**Example 1:** If  $y = \left( \sqrt{\frac{1+\cos 2x}{1-\cos 2x}} + \sqrt{\sec^2 x - 1} \right)^{-1} + (1+x)^n$ , then show that  $\frac{dy}{dx} - n(1+x)^{n-1} = \cos 2x$ .

**Solution:**

We have

$$\begin{aligned}
 y &= \left( \sqrt{\frac{1+\cos 2x}{1-\cos 2x}} + \sqrt{\sec^2 x - 1} \right)^{-1} + (1+x)^n \\
 &= \left( \sqrt{\frac{\cos^2 x}{\sin^2 x}} + \sqrt{\tan^2 x} \right)^{-1} + \sum_{r=0}^n {}^n C_r x^r \\
 &= (\cot x + \tan x)^{-1} + \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right) \\
 &= \left( \frac{\cos x}{\sin x} + \frac{\sin x}{\cos x} \right)^{-1} + \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right) \\
 &= \left( \frac{\sin^2 x + \cos^2 x}{\sin x \cos x} \right)^{-1} + \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right) \\
 &= \sin x \cdot \cos x + \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right)
 \end{aligned}$$

Hence,

$$\frac{dy}{dx} = \frac{d}{dx} (\sin x \cdot \cos x) + \frac{d}{dx} \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right)$$

Now,

$$\begin{aligned}
 \frac{d}{dx} (\sin x \cdot \cos x) &= \frac{d}{dx} (\sin x) \cdot \cos x + \sin x \cdot \frac{d}{dx} (\cos x) \\
 &= \cos x \cdot \cos x + \sin x (-\sin x) \\
 &= \cos^2 x - \sin^2 x \\
 &= \cos 2x
 \end{aligned}$$

$$\begin{aligned}
& \frac{d}{dx} \left( 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 \dots + \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} + \frac{n(n-1)\dots 1}{n!} x^n \right) \\
&= \frac{d}{dx} (1) + \frac{d}{dx} (nx) + \frac{d}{dx} \left( \frac{n(n-1)}{2!} x^2 \right) + \frac{d}{dx} \left( \frac{n(n-1)(n-2)}{3!} x^3 \right) \dots + \frac{d}{dx} \left( \frac{n(n-1)\dots 2}{(n-1)!} x^{n-1} \right) + \frac{d}{dx} \left( \frac{n(n-1)\dots 1}{n!} x^n \right) \\
&= 0 + n \frac{d}{dx} (x) + \frac{n(n-1)}{2!} \frac{d}{dx} (x^2) + \frac{n(n-1)(n-2)}{3!} \frac{d}{dx} (x^3) + \dots + \frac{n(n-1)\dots 2}{(n-1)!} \frac{d}{dx} (x^{n-1}) + \frac{n(n-1)\dots 1}{n!} \frac{d}{dx} (x^n) \\
&= n + \frac{2n(n-1)}{2!} x + \frac{3n(n-1)(n-2)}{3!} x^2 + \dots + \frac{(n-1)n(n-1)\dots 2}{(n-1)!} (x^{n-2}) + \frac{n(n-1)\dots 1}{n!} (x^{n-1}) \\
&= n \left( 1 + (n-1)x + \frac{(n-1)(n-2)}{2!} x^2 + \dots + \frac{(n-1)(n-2)\dots 2}{(n-2)!} x^{n-2} + \frac{(n-1)(n-2)\dots 1}{(n-1)!} x^{n-1} \right) \\
&= n(1+x)^{n-1}
\end{aligned}$$

Hence,

$$\begin{aligned}
\frac{dy}{dx} &= \cos 2x + n(1+x)^n \\
\Rightarrow \cos 2x &= \frac{dy}{dx} - n(1+x)^n
\end{aligned}$$

**Example 2:** Find  $\frac{dy}{dx}$  if  $y = \frac{2x^7 + 3 + \tan x}{x(\sin x - \cos x)}$ .

**Solution:**

$$\begin{aligned}
y &= \frac{2x^7 + 3 + \tan x}{x(\sin x - \cos x)} = \frac{2x^7 + 3 + \tan x}{(x \sin x - x \cos x)} \\
\Rightarrow \frac{dy}{dx} &= \frac{(2x^7 + 3 + \tan x)' \cdot (x \sin x - x \cos x) - (2x^7 + 3 + \tan x) \cdot (x \sin x - x \cos x)'}{(x \sin x - x \cos x)^2} \quad \dots (1) \\
&\quad \because \left( \frac{u}{v} \right)' = \frac{u'v - uv'}{v^2}
\end{aligned}$$

Now,

$$\begin{aligned}
(2x^7 + 3 + \tan x)' &= \frac{d}{dx}(2x^7 + 3 + \tan x) \\
&= 2 \frac{d}{dx}(x^7) + \frac{d}{dx}(3) + \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) \\
&= 2 \times 7x^6 + 0 + \frac{\frac{d}{dx}(\sin x) \cdot \cos x - \sin x \cdot \frac{d}{dx}(\cos x)}{\cos^2 x} \\
&= 14x^6 + \frac{\cos x \cdot \cos x - \sin x \cdot (-\sin x)}{\cos^2 x} \\
&= 14x^6 + \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\
&= 14x^6 + \sec^2 x
\end{aligned}$$

$$\begin{aligned}
(x \sin x - x \cos x)' &= \frac{d}{dx}(x \sin x - x \cos x) \\
&= \frac{d}{dx}(x \sin x) - \frac{d}{dx}(x \cos x) \\
&= \frac{d}{dx}(x) \cdot (\sin x) + x \cdot \frac{d}{dx}(\sin x) - \left( \frac{d}{dx}(x) \cdot (\cos x) + x \cdot \frac{d}{dx}(\cos x) \right) \quad [(uv)' = u'v + uv'] \\
&= \sin x + x \cos x - [\cos x + x(-\sin x)] \\
&= (1 + x) \sin x + (x - 1) \cos x
\end{aligned}$$

On substituting all the values in equation (1), we obtain

$$\begin{aligned}
\frac{dy}{dx} &= \frac{(2x^7 + 3 + \tan x)' \cdot (x \sin x - x \cos x) - (2x^7 + 3 + \tan x) \cdot (x \sin x - x \cos x)'}{(x \sin x - x \cos x)^2} \\
&= \frac{(14x^6 + \sec^2 x) \cdot (x \sin x - x \cos x) - (2x^7 + 3 + \tan x) \cdot [(1 + x) \sin x + (x - 1) \cos x]}{x^2 (\sin x - \cos x)^2}
\end{aligned}$$